

little, if any, at Buffalo, N. Y. The western margins of the storm were near Sandusky, without damage.

Storm-sewer construction and catch basins in the areas immediately near Public Square appeared to be adequate for the run-off of this storm, but in the highlands and eastern suburbs, either the recent construction of water-courses and their resultant constriction is woefully undersized, or else the rain in that region was much greater than down town.

The loss, according to newspaper headlines, was in the millions.

A TORNADO IN NEW MEXICO¹

By C. E. LINNEY

[Weather Bureau Office, Santa Fe, N. Mex.]

Just about a year after a destructive tornado struck Wagon Mound, Mora County, N. Mex., a second tornadic

storm was observed to form on June 5, 1931, near the small village of French, Colfax County, about 35 miles northeast of Wagon Mound. This second tornado moved slowly east-southeast through a thinly settled country, doing but relatively little damage by reason of the sparsely settled country. It passed into and across Union County, doing quite a bit of damage to buildings and causing the death of a 3-year-old girl by an out-building crashing upon her. The property loss in the Gladstone district is estimated at about \$1,500 and in the Barney and Sedan districts about \$10,000 and \$20,000, respectively.

The tornado was under observation during its entire course of about 90 miles. It dissipated after crossing the Texas border.

¹ Condensed from the original.—Ed.

BIBLIOGRAPHY

C. FITZHUGH TALMAN, in charge of Library

RECENT ADDITIONS

The following have been selected from among the titles of books recently received as representing those most likely to be useful to Weather Bureau officials in their meteorological work and studies:

Bjerknes, V., & Solberg, H.

Zellulare Trägheitswellen und Turbulenz. Oslo. 1929. 16 p. 24 cm. (Norske videnskaps-akad. i Oslo i matem.-natur. kl. 1929. no. 7.)

Bruzon, E., & Carton, P.

Le climat de l'Indochine et les typhons de la mer de Chine. Hanoi. 1930. 310 p. figs. plates (part fold.). 27½ cm. (Num. spec. des Annales du serv. met.)

Franssila, M.

Die Häufigkeit der verschiedenen Windgeschwindigkeiten am aerologischen Observatorium Ilmala. Helsingfors. 1930. 15 p. figs. 24 cm. (Mitt. des Met. inst. der univ., Helsingfors. N:o 16.)

Gamba, Pericle.

Sulla temperatura del suolo a Pavia. (Risultati di un ventennio di osservazioni geotermiche nel R. Osservatorio geofisico.) Roma. 1929. 50 p. plates (part fold.). 25 cm. (Parma. Uff. idrog. del Po. Pubb. n. 11.)

Great Britain. Air ministry. Airship services division.

Conditions along airship routes. Preliminary report on routes between England (Cardington), Egypt (Ismailia), and India (Karachi). September. n. p. n. d. 24 p. charts (fold.). 33 cm. (Airship met'l rep. 80.)

International commission for the exploration of the upper air.

Ergebnisse der aerologischen Messungen. 24. I. 1925: 17–22. VIII. 1925. Berlin. 1930. 23, 172 p. 30 cm. [Author, title and text in German, French, and English.]

Jenny, Hans.

Study on the influence of climate upon the nitrogen and organic matter content of the soil. Columbia. 1930. 66 p. figs. 23 cm. (Univ. Mo. Coll. agric. Agr. exper. sta. res. bull. 152.)

Johansson, Osc. V.

Der mittlere Verlauf der Lufttemperatur in der Winternacht. Helsingfors. 1931. 26 p. 24 cm. (Mitt. des met. Inst. der Univ., Helsingfors. N:o 17.)

Middleton, W. E. Knowles.

Measurement of visibility at night. p. 39–48. figs. 25 cm. (Trans. R. S. C. sec. III, 1931.)

Miller, A. Austin.

Climatology. London. [1931.] x, 304 p. figs. 23 cm.

Molchanov, P. A.

Investigation of upper layers of atmosphere by method of radio sounding in Arctic condition. Leningrad. 1931. 12 p. figs. 28 cm. (Repr.: Mag. sci. res. bur. 1st issue.) [Author, title and text in Russian. English translation typewritten.]

Nagler, Floyd A.

Drought of 1930 in Iowa. Iowa City. 1931. 16 p. figs. 23½ cm. (Univ. Iowa exten. bull. Bull. no. 266. Apr. 15, 1931.)

Petersen, Helge.

Theoretical determination of the heights of the stratosphere, the ozone layer and the height of maximum luminosity of the aurora. København. 1931. 19 p. figs. 25 cm. (Pub. Danske met. inst. Commun. mag. no. 13.)

Pettersson, O.

Flödskrifter och vattenutbytet mellan de tropiska och de Polara haven. En studie i geofysik och kosmisk fysik. [Göteborg. 1930.] 18, 10 p. illus. 34 cm. (Svenska hydrog.-biol. komm. skrifter. Ny ser. Hydrografi VIII.)

Schultze, Herbert.

Die Schwankungen der barometrischen Hochlagen über Mitteleuropa und ihr Einfluss auf die Ernterträge in Deutschland. Halle. 1928. 141 p. figs. 25 cm. (Inaug.-Dissert. Verein. Friedrichs-Univ. Halle-Wittenberg.)

Svensson, Aron.

Verdunstung und Abkühlung oder Erwärmung in einem laminaren Gasstrom von konstanter Geschwindigkeit. Anwendung auf Assmann's Aspirations-psychrometer. Stockholm. 1931. 28 p. figs. 22 cm. (Arkiv för mat., astron. och fysik utg. av K. Svenska vetenskapsakad. Bd. 22 A. N:o 23.)

Taylor, C. Fayette.

Suggested method for measuring turbulence. Washington. 1931. 7 p. plates. 26½ cm. (Tech. notes, Nat. adv. comm. for aeron. No. 380.) [Manifolded.]

Väistälä, Vilho.

Die Wolkenhöhen in Finnland. Helsingfors. 1930. 28 p. figs. 24½ cm. (Mitt. des met. Inst. der Univ., Helsingfors. N:o 15.)

Waring, F. H., & Stewart, F. D.

Effect of the 1930 drought upon Ohio public water supplies. p. 18–23. 23 cm. (Engin. exper. sta. news. Ohio state univ. Suppl. v. 3, no. 3. June, 1931.)

Welter, L.

La pluie à Dakar et l'activité solaire. p. 264–271. figs. 25½ cm. (Bull. du com. d'études hist. et sci. de l'Afrique occid. franc. T. 13, no. 2. Année 1930. Avr.-juin.)